Contents

Volume 105 1993

Nakata, T., Takemura, R. and Hirokawa, N.
A novel member of the dynamin family of GTP-binding proteins is expressed specifically in the testis 1-5

Gonzalez-Fernandez, F., Kittredge, K. L., Rayborn, M. E., Hollyfield, J. G., Landers, R. A., Saha, M. and Grainger, R. M.
Interphotoreceptor retinoid-binding protein (IRBP), a major 124 kDa glycoprotein in the interphotoreceptor matrix of *Xenopus laevis*: characterization, molecular cloning and biosynthesis 7-22

Hashino, E. and Salvi, R. J.
Changing spatial patterns of DNA replication in the noise-damaged chick cochlea 23-31

Thiry, M.
Ultrastructural distribution of DNA and RNA within the nucleolus of human Sertoli cells as seen by molecular immunocytochemistry 33-39

Carmena, M., Abad, J. P., Villasante, A. and Gonzalez, C.
The *Drosophila melanogaster dodecasatellite* sequence is closely linked to the centromere and can form connections between sister chromatids during mitosis 41-50

Cross, D., Vial, C. and Maccioni, R. B.
A tau-like protein interacts with stress fibers and microtubules in human and rodent cultured cell lines 51-60

Shankar, G., Davison, I., Helfrich, M. H., Mason, W. T. and Horton, M. A.
Integrin receptor-mediated mobilisation of intranuclear calcium in rat osteoclasts 61-68

Baptist, M., Dumont, J. E. and Roger, P. P.
Demonstration of cell cycle kinetics in thyroid primary culture by immunostaining of proliferating cell nuclear antigen: differences in cyclic AMP-dependent and -independent mitogenic stimulations 69-80

Fredieu, J. R. and Mahowald, A. P.
Characterization of a putative *Drosophila* GTP-binding protein 81-91

Mullett, H., Ratcliffe, N. A. and Rowley, A. F.
The generation and characterisation of anti-insect blood cell monclonal antibodies 93-100

Pasqualini, R., Bodorova, J., Ye, S. and Hemler, M. E.
A study of the structure, function and distribution of β_5 integrins using novel anti-β_5 monoclonal antibodies 101-111

Adolph, S., Brüsselbach, S. and Müller, R.
Inhibition of transcription blocks cell cycle progression of NIH3T3 fibroblasts specifically in G_1 113-122

Lucibello, F. C., Sewing, A., Brüsselbach, S., Bürger, C. and Müller, R.
Deregulation of cyclins D1 and E and suppression of cdk2 and cdk4 in senescent human fibroblasts 123-133

Samejima, I., Matsumoto, T., Nakaseko, Y., Beach, D. and Yanagida, M.
Identification of seven new *cut* genes involed in *Schizosaccharomyces pombe* mitosis 135-143

Julian, M., Tollon, Y., Lajoie-Mazenc, I., Moisand, A., Mazarguil, H., Puget, A. and Wright, M.
γ-Tubulin participates in the formation of the midbody during cytokinesis in mammalian cells 145-156

Gueth-Hallonet, C., Antony, C., Aghion, J., Santa-Maria, A., Lajoie-Mazenc, I., Wright, M. and Maro, B.
γ-Tubulin is present in acentriolar MTOCs during early mouse development 157-166

Lee, T.-L., Lin, Y.-C., Mochitate, K. and Grinnell, F.
Stress-relaxation of fibroblasts in collagen matrices triggers ectocytosis of plasma membrane vesicles containing actin, annexins II and VI, and β_1 integrin receptors 167-177

Zambruno, G., Marchisio, P. C., Melchiori, A., Bondanza, S., Cancedda, R. and De Luca, M.
Expression of integrin receptors and their role in adhesion, spreading and migration of normal human melanocytes 179-190

Thomas, L., Chan, P. W., Chang, S. and Damsky, C.
5-Bromo-2-deoxyuridine regulates invasiveness and expression of integrins and matrix-degrading proteinases in a differentiated hamster melanoma cell 191-201

Clark, P., Britland, S. and Connolly, P.
Growth cone guidance and neuron morphology on micropatterned laminin surfaces 203-212

Rooney, P., Wang, M., Kumar, P. and Kumar, S.
Angiogenic oligosaccharides of hyaluronan enhance the production of collagens by endothelial cells 213-218

Franck, Z., Gary, R. and Bretscher, A.
Meosin, like ezrin, colocalizes with actin in the cortical cytoskeleton in cultured cells, but its expression is more variable 219-231

Yumura, S. and Kitanishi-Yumura, T.
A mechanism for the intracellular localization of myosin II filaments in the *Dictyostelium* amoeba 233-242

Fuchs, M., Jones, M. K. and Williams, K. L.
Characterisation of an epithelium-like layer of cells in the multicellular *Dictyostelium discoideum* slug 243-253

Flaadt, H., Jaworski, E., Schlatterer, C. and Malchow, D.
Cyclic AMP- and Ins(1,4,5)P_3-induced Ca^{2+} fluxes in permeabilised cells of *Dictyostelium discoideum*: cGMP regulates Ca^{2+} entry across the plasma membrane 255-261

Kurkdjian, A., Leitz, G., Manigault, P., Harim, A. and Greulich, K. O.
Non-enzymatic access to the plasma membrane of *Medicago* root hairs by laser microsurgery 263-268

Contents

Commentary

Scott, P. A. E. and Bicknell, R.
The isolation and culture of microvascular endothelium 269-273

Hashimoto, T., Amagai, M., Parry, D. A. D., Dixon, T. W., Tsukita, S., Tsukita, Sh., Miki, K., Sakai, K., Inokuchi, Y., Kudoh, J., Schimizu, N. and Nishikawa, T.
Desmoyokin, a 680 kDa keratinocyte plasma membrane-associated protein, is homologous to the protein encoded by human gene AHNAK 275-286

Baumann, O. and Takeyasu, K.
Polarized distribution of Na,K-ATPase in honeybee photoreceptors is maintained by interaction with glial cells 287-302

Wu, R.-L., Galvin, S., Wu, S.-K., Xu, C., Blumenberg, M. and Sun, T.-T.
A 300 bp 5′-upstream sequence of a differentiation-dependent rabbit K3 keratin gene can serve as a keratinocyte-specific promoter 303-316

Guido, S. and Tranquillo, R. T.
A methodology for the systematic and quantitative study of cell contact guidance in oriented collagen gels. Correlation of fibroblast orientation and gel birefringence 317-331

Rajasekaran, A. K., Morimoto, T., Hanzel, D. K., Rodriguez-Boulan, E. and Kreibich, G.
Structural reorganization of the rough endoplasmic reticulum without size expansion accounts for dexamethasone-induced secretory activity in AR42J cells 333-345

Sahlas, D. J., Milankov, K., Park, P. C. and De Boni, U.
Distribution of snRNPs, splicing factor SC-35 and actin in interphase nuclei: immunocytochemical evidence for differential distribution during changes in functional states 347-357

Ouspenski, I. I. and Brinkley, B. R.
Centromeric DNA cloned from functional kinetochore fragments in mitotic cells with unreplicated genomes 359-367

Akins, R. E. and Tuan, R. S.
Transepithelial calcium transport in the chick chorioallantoic membrane. I. Isolation and characterization of chorionic ectoderm cells 369-379

Akins, R. E. and Tuan, R. S.
Transepithelial calcium transport in the chick chorioallantoic membrane. II. Compartmentalization of calcium during uptake 381-388

Li, X. and Etkin, L. D.
Xenopus nuclear factor 7 (xnf7) possesses an NLS that functions efficiently in both oocytes and embryos 389-395

Braga, V. M. M. and Gendler, S. J.
Modulation of Muc-1 mucin expression in the mouse uterus during the estrus cycle, early pregnancy and placentation 397-404

Flouriot, G., Vaillant, C., Salbert, G., Pelissero, C., Guiraud, J. M. and Valotaire, Y.
Monolayer and aggregate cultures of rainbow trout hepatocytes: long-term and stable liver-specific expression in aggregates 407-416

Chambers, J. A. and Harris, A.
Expression of the cystic fibrosis gene and the major pancreatic mucin gene, MUC1, in human ductal epithelial cells 417-422

Griffin, M., Bhandari, R., Hamilton, G., Chan, Y.-C. and Powell, J. T.
Alveolar type II cell-fibroblast interactions, synthesis and secretion of surfactant and type I collagen 423-432

Chou, C.-F., Riopel, C. L., Rott, L. S. and Omary M. B.
A significant soluble keratin fraction in 'simple' epithelial cells. Lack of an apparent phosphorylation and glycosylation role in keratin solubility 433-444

Peterson, M. D. and Mooseker, M. S.
An in vitro model for the analysis of intestinal brush border assembly. I. Ultrastructural analysis of cell contact-induced brush border assembly in Caco-2_{BBe} cells 445-460

Peterson, M. D., Bement, W. M. and Mooseker, M. S.
An in vitro model for the analysis of intestinal brush border assembly. II. Changes in expression and localization of brush border proteins during cell contact-induced brush border assembly in Caco-2_{BBe} cells 461-472

Lawrence, B. P. and Brown, W. J.
Inhibition of protein synthesis separates autophagic sequestration from the delivery of lysosomal enzymes 473-480

Frangioni, J. V. and Neel, B. G.
Use of a general purpose mammalian expression vector for studying intracellular protein targeting: identification of critical residues in the nuclear lamin A/C nuclear localization signal 481-488

Dunlevy, J. R. and Couchman, J. R.
Controlled induction of focal adhesion disassembly and migration in primary fibroblasts 489-500

Faassen, A. E., Mooradian, D. L., Tranquillo, R. T., Dickinson, R. B., Letourneau, P. C., Oegema, T. R. and McCarthy, J. B.
Cell surface CD44-related chondroitin sulfate proteoglycan is required for transforming growth factor-β-stimulated mouse melanoma cell motility and invasive behavior on type 1 collagen 501-511

Bohrmann, J.
Antisera against a channel-forming 16 kDa protein inhibit dye-coupling and bind to cell membranes in *Drosophila* ovarian follicles 513-518

Boschelli, F., Uptain, S. M. and Lightbody, J. J.
The lethality of $p60^{v\text{-}src}$ in *Saccharomyces cerevisiae* and activation of $p34^{CDC28}$ kinase are dependent on the integrity of the SH2 domain 519-528

Novak, B., Sveiczer, A. and Mitchison, J. M.
CO_2 production in cell-free extracts of fission yeast detects cell cycle changes 529-531

Vishwanatha, J. K. and Kumble, S.
Involvement of annexin II in DNA replication: evidence from cell-free extracts of *Xenopus* eggs 533-540

Hassan, A. B. and Cook, P. R.
Visualization of replication sites in unfixed human cells 541-550

Giannasca, P. J., Horowitz, R. A. and Woodcock, C. L.
Transitions between in situ and isolated chromatin 551-561

Clarke, D. J., Johnson, R. T. and Downes, C. S.
Topoisomerase II inhibition prevents anaphase chromatid segregation in mammalian cells independently of the generation of DNA strand breaks 563-569

Czaban, B. B., Forer, A. and Bajer, A. S.
Ultraviolet microbeam irradiation of chromosomal spindle fibres in *Haemanthus katherinae* endosperm. I. Behaviour of the irradiated region 571-578

Lin, S. X. H. and Collins, C. A.
Regulation of the intracellular distribution of cytoplasmic dynein by serum factors and calcium 579-588

Maekawa, T. and Kuriyama, R.
Primary structure and microtubule-interacting domain of the SP-H antigen: a mitotic map located at the spindle pole and characterized as a homologous protein to NuMA 589-600

Commentary

Screpanti, I., Modesti, A. and Gulino, A.
Heterogeneity of thymic stromal cells and thymocyte differentiation: a cell culture approach 601-606

Vojtěšek, B. and Lane, D. P.
Regulation of p53 protein expression in human breast cancer cell lines 607-612

David-Pfeuty, T., Bagrodia, S. and Shalloway, D.
Differential localization patterns of myristoylated and nonmyristoylated c-Src proteins in interphase and mitotic c-Src overexpresser cells 613-628

Rebut-Bonneton, C., Boutemy-Roulier, S. and Evain-Brion, D.
Modulation of $pp60^{c\text{-}src}$ activity and cellular localization during differentiation of human trophoblast cells in culture 629-636

Turner, C. E., Schaller, M. D. and Parsons, J. T.
Tyrosine phosphorylation of the focal adhesion kinase $pp125^{FAK}$ during development: relation to paxillin 637-645

Meriläinen, J., Palovuori, R., Sormunen, R., Wasenius, V.-M. and Lehto, V.-P.
Binding of the α-fodrin SH3 domain to the leading lamellae of the locomoting chicken fibroblasts 647-654

Che, A., Cherry, R. J., Bannister, L. H. and Dluzewski, A. R.
Aggregation of band 3 in hereditary ovalocytic red blood cell membranes. Electron microscopy and protein rotational diffusion studies 655-660

Cormier, F.
Avian pluripotent haemopoietic progenitor cells: detection and enrichment from the para-aortic region of the early embryo 661-666

du Cros, D. L., Isaacs, K. and Moore, G. P. M.
Distribution of acidic and basic fibroblast growth factors in ovine skin during follicle morphogenesis 667-674

Zaffran, Y., Lepidi, H., Bongrand, P., Mege, J.-L. and Capo, C.
F-actin content and spatial distribution in resting and chemoattractant-stimulated human polymorphonuclear leucocytes. Which role for intracellular free calcium? 675-684

Blencowe, B. J., Carmo-Fonseca, M., Behrens, S.-E., Lurhmann, R. and Lamond, A. I.
Interaction of the human autoantigen p150 with splicing snRNPs 685-697

Kocks, C., Hellio, R., Gounon, P., Ohayon, H. and Cossart, P.
Polarized distribution of *Listeria monocytogenes* surface protein ActA at the site of directional actin assembly 699-710

Maldonado-Codina, G., Llamazares, S. and Glover, D. M.
Heat shock results in cell cycle delay and synchronisation of mitotic domains in cellularised *Drosophila melanogaster* embryos 711-720

Buttgereit, D.
Redundant enhancer elements guide β1 tubulin gene expression in apodemes during *Drosophila* embryogenesis 721-727

Georgieff, I. S., Liem, R. K. H., Couchie, D., Mavilia, C., Nunez, J. and Shelanski, M. L.
Expression of high molecular weight tau in the central and peripheral nervous systems 729-737

Grenz, H., Carbonetto, S. and Goodman, S. L.
$\alpha_3\beta_1$ Integrin is moved into focal contacts in kidney mesangial cells 739-751

Langhofer, M., Hopkinson, S. B. and Jones, J. C. R.
The matrix secreted by 804G cells contains laminin-related components that participate in hemidesmosome assembly in vitro 753-764

Friederich, E., Kreis, T. E. and Louvard, D.
Villin-induced growth of microvilli is reversibly inhibited by cytochalasin D 765-775

Vojtek, A. B. and Cooper, J. A.
Identification and characterization of a cDNA encoding mouse CAP: a homolog of the yeast adenylyl cyclase associated protein 777-785

Azarian, S. M., Schlamp, C. L. and Williams, D. S.
Characterization of calpain II in the retina and photoreceptor outer segments 787-798

Schmidt-Zachmann, M. S. and Nigg, E. A.
Protein localization to the nucleolus: a search for targeting domains in nucleolin 799-806

Mah, D. C. W., Dijkwel, P. A., Todd, A., Klein, V., Price, G. B. and Zannis-Hadjopoulos, M.
ors12, a mammalian autonomously replicating DNA sequence, associates with the nuclear matrix in a cell cycle-dependent manner 807-818

Egea, G., Francí, C., Gambús, G., Lesuffleur, T., Zweibaum, A. and Real, F. X.
cis-Golgi resident proteins and *O*-glycans are abnormally compartmentalized in the RER of colon cancer cells 819-830

Contents

Beghdadi-Rais, C., Schreyer, M., Rousseaux, M., Borel, P., Eisenberg, R. J., Cohen, G. H., Bron, C. and Fasel, N.
Carboxyl terminus structural requirements for glycosyl-phosphatidylinositol anchor addition to cell surface proteins 831-840

Kensler, R. W. and Stewart, M.
The relaxed crossbridge pattern in isolated rabbit psoas muscle thick filaments 841-848

Nolta, K. V., Padh, H. and Steck, T. L.
An immunocytochemical analysis of the vacuolar protein pump in *Dictyostelium discoideum* 849-859

Aubry, L., Klein, G., Martiel, J.-L. and Satre, M.
Kinetics of endosomal pH evolution in *Dictyostelium discoideum* amoebae. Study by fluorescence spectroscopy 861-866

Commentary

Fulton, A. B. and L'Ecuyer, T.
Cotranslational assembly of some cytoskeletal proteins: implications and prospects 867-871

Ookata, K., Hisanaga, S.-i., Okumura, E. and Kishimoto, T.
Association of $p34^{cdc2}$/cyclin B complex with microtubules in starfish oocytes 873-881

Hosoya, N., Mitsui, M., Yazama, F., Ishihara, H., Ozaki, H., Karaki, H., Hartshorne, D. J. and Mohri, H.
Changes in the cytoskeletal structure of cultured smooth muscle cells induced by calyculin-A 883-890

Jiang, C.-J. and Sonobe, S.
Identification and preliminary characterization of a 65 kDa higher-plant microtubule-associated protein 891-901

Triviños-Lagos, L., Ohmachi, T., Albrightson, C., Burns, R. G., Ennis, H. L. and Chisholm, R. L.
The highly divergent α- and β-tubulins from *Dicytostelium discoideum* are encoded by single genes 903-911

Cheek, T. R., Morgan, A., O'Sullivan, A. J., Moreton, R. B., Berridge, M. J. and Burgoyne, R. D.
Spatial localization of agonist-induced Ca^{2+} entry in bovine adrenal chromaffin cells. Different patterns induced by histamine and angiotensin II, and relationship to catecholamine release 913-921

Navarro, P., Lozano, E. and Cano, A.
Expression of E- or P-cadherin is not sufficient to modify the morphology and the tumorigenic behavior of murine spindle carcinoma cells. Possible involvement of plakoglobin 923-934

Tixier-Vidal, A., Barret, A., Picart, R., Mayau, V., Vogt, D., Wiedenmann, B. and Goud, B.
The small GTP-binding protein, Rab6p, is associated with both Golgi and post-Golgi synaptophysin-containing membranes during synaptogenesis of hypothalamic neurons in culture 935-947

Loveridge, N., Farquharson, C., Hesketh, J. E., Jakowlew, S. B., Whitehead, C. C. and Thorp, B. H.
The control of chondrocyte differentiation during endochondral bone growth in vivo: changes in TGF-β and the proto-oncogene c-*myc* 949-956

Yao, S.-N. and Kurachi, K.
Implanted myoblasts not only fuse with myofibers but also survive as muscle precursor cells 957-963

Witmer-Pack, M. D., Crowley, M. T., Inaba, K. and Steinman, R. M.
Macrophages, but not dendritic cells, accumulate colloidal carbon following administration in situ 965-973

Wardale, R. J. and Duance, V. C.
Quantification and immunolocalisation of porcine articular and growth plate cartilage collagens 975-984

Gourdie, R. G., Severs, N. J., Green, C. R., Rothery, S., Germroth, P. and Thompson, R. P.
The spatial distribution and relative abundance of gap-junctional connexin40 and connexin43 correlate to functional properties of components of the cardiac atrioventricular conduction system 985-991

Peifer, M.
The product of the *Drosophila* segment polarity gene *armadillo* is part of a multi-protein complex resembling the vertebrate adherens junction 993-1000

Sriramarao, P., Mendler, M. and Bourdon, M. A.
Endothelial cell attachment and spreading on human tenascin is mediated by $\alpha_2\beta_1$ and $\alpha_v\beta_3$ integrins 1001-1012

Montesano, R., Pepper, M. S. and Orci, L.
Paracrine induction of angiogenesis in vitro by Swiss 3T3 fibroblasts 1013-1024

Berryman, M., Franck, Z. and Bretscher, A.
Ezrin is concentrated in the apical microvilli of a wide variety of epithelial cells whereas moesin is found primarily in endothelial cells 1025-1043

Bard, J. B. L., Hulmes, D. J. S., Purdom, I. F. and Ross, A. S. A.
Chick corneal development in vitro: diverse effects of pH on collagen assembly 1045-1055

Remington, S. G.
Chicken filensin: a lens fiber cell protein that exhibits sequence similarity to intermediate filament proteins 1057-1068

Mitchell, D. R. and Kang, Y.
Reversion analysis of dynein intermediate chain function 1069-1078

De Luca, M., Siegrist, W., Bondanza, S., Mathor, M., Cancedda, R. and Eberle, A. N.
α Melanocyte stimulating hormone (αMSH) stimulates normal human melanocyte growth by binding to high-affinity receptors 1079-1084

Reiland, J. and Rapraeger, A. C.
Heparan sulfate proteoglycan and FGF receptor target basic FGF to different intracellular destinations 1085-1093

Byrne, S. M. and Hoffman, C. S.
Six *git* genes encode a glucose-induced adenylate cyclase activation pathway in the fission yeast *Schizosaccharomyces pombe* 1095-1100

Bangs, J. D., Uyetake, L., Brickman, M. J., Balber, A. E. and Boothroyd, J. C.
Molecular cloning and cellular localization of a BiP homologue in *Trypanosoma brucei*. Divergent ER retention signals in a lower eukaryote 1101-1113

Pidoux, A. L. and Armstrong, J.
The BiP protein and the endoplasmic reticulum of *Schizosaccharomyces pombe*: fate of the nuclear envelope during cell division 1115-1120

Lang, S., Decristoforo, T., Waitz, W. and Loidl, P.
Biochemical and morphological characterization of the nuclear matrix during the synchronous cell cycle of *Physarum polycephalum* 1121-1130

Flaadt, H., Jaworski, E. and Malchow, D.
Evidence for two intracellular calcium pools in *Dictyostelium*: the cAMP-induced calcium influx is directed into a NBD-Cl- and 2,5-di-(tert-butyl)-1,4-hydroquinone-sensitive pool 1131-1135

Morgans, C. W. and Kopito, R. R.
Association of the brain anion exchanger, AE3, with the repeat domain of ankyrin 1137-1142

Jackson, D. A. and Cook, P. R.
Transcriptionally active minichromosomes are attached transiently in nuclei through transcription units 1143-1150

Highett, M. I., Beven, A. F. and Shaw, P. J.
Localization of 5 S genes and transcripts in *Pisum sativum* nuclei 1151-1158

Hayashi-Isimaru, Y., Ueda, K. and Nonaka, M.
Detection of DNA in the nucleoids of chloroplasts and mitochondria in *Euglena gracilis* by immunoelectron microscopy 1159-1164

García de Herreros, A., Fabre, M., Batlle, E., Balagué, C. and Real, F. X.
The tumor promoter 12-*O*-tetradecanoylphorbol-13-acetate blocks differentiation of HT-29 human colon cancer cells 1165-1172

Shepherd, V. A., Orlovich, D. A. and Ashford, A. E.
Cell-to-cell transport via motile tubules in growing hyphae of a fungus 1173-1178

Index of Authors and Titles 1179-1184

Subject Index 1185-1197